

Example Client, XYZ123 1234 Warde Road Ann Arbor MI 48108 EXAMPLE, REPORT W WX0000003826 F 12/05/1988 34 Y

Referral Testing							
		Collec	cted: 09/18/2023	3 15:27	Received:	09/18/2023	15:27
Test Nam	2	<u>Result</u>	<u>Flag</u>	Ref-Range	es	<u>Units</u>	<u>Site</u>
Methylphenidate and Metabolite, Serum/PlasmaMethylphenidate32Hng/mLNMF							NMRL
	Reporting Limit: 2.0 ng/mL Synonym(s): Ritalin(R) Peak plasma concentrations of 8-22 ng/mL are usual at 1 to 2 hours following 10 to 20 mg oral pediatric anti-hyperkinetic doses. (Methylphenidate specimens should be kept frozen). Analysis by High Performance Liquid Chromatography/ Tandem Mass Spectrometry (LC-MS/MS)						
Ritalinic A	cid	662	н			ng/mL	NMRL
Reporting Limit: 10 ng/mL Synonym(s): Methylphenidate Metabolite Plasma concentrations 3 to 6 hours post-dose in children given a 10 to 15 mg oral dose of Methylphenidate: 80-250 ng Ritalinic Acid/mL. Analysis by High Performance Liquid Chromatography/ Tandem Mass Spectrometry (LC-MS/MS) This test was developed and its performance characteristics determined by NMS Labs. It has not been cleared or approved by the US Food and Drug Administration. Testing performed at NMS Labs, Inc. 200 Welsh Road Horsham, PA 19044-2208							
	CLIA 39D0197898						
			Reported	I Date: 20	23.09.18	15:27 MF	PDR

LAB: L - LOW, H - HIGH, AB - ABNORMAL, C - CRITICAL, . - NOT TESTED

Ordered By: KAJAL SITWALA, MD, PhD WX0000000002353