



LABORATORY REPORT

Example Client, XYZ123
1234 Warde Road
Ann Arbor MI 48108

EXAMPLE, REPORT
WX0000003039 M 12/05/1988 34 Y

Immunochemistry

Collected: 04/29/2022 07:22 Received: 04/29/2022 07:22

Table with 6 columns: Test Name, Result, Flag, Ref-Ranges, Units, Site. Row 1: Thyroglobulin and Anti-Thyroglobulin Antibody Panel. Row 2: Anti-thyroglobulin Antibody, 23, H, <4, IU/mL, WMRL.

According to the reagent manufacturer, consumption of biotin supplements, or multivitamins containing biotin, may interfere with the results of this assay. For individuals taking biotin-containing supplements, testing at least three days after cessation of supplement consumption is recommended.

Table with 6 columns: Test Name, Result, Flag, Ref-Ranges, Units, Site. Row 1: Thyroglobulin, 44.0, SeeBelow, ng/mL, WMRL.

Thyroglobulin Reference Range:
Intact thyroid: 1.6-50.0 ng/mL
Athyrotic, post-thyroidectomy (tumor marker): <0.1 ng/mL

Thyroglobulin antibodies can interfere with the determination of thyroglobulin. If the specimen contains thyroglobulin antibodies, please interpret the thyroglobulin result with caution.

The Beckman Access DXI chemiluminescent immunoassay is used. Results obtained with different assay methods or kits cannot be used interchangeably. Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.

According to the reagent manufacturer, consumption of biotin supplements, or multivitamins containing biotin, may interfere with the results of this assay. For individuals taking biotin-containing supplements, testing at least three days after cessation of supplement consumption is recommended.

Reported Date: 04/29/2022 07:22 THY

Performing Site: WMRL: WARDE MEDICAL LABORATORY 300 West Textile Road Ann Arbor MI 48108

LAB: L - LOW, H - HIGH, AB - ABNORMAL, C - CRITICAL, . - NOT TESTED

D629000000 Ordered By: CLIENT CLIENT
WX0000003039 WX00000000001595
Printed D&T: 12/01/23 11:10

Kajal V. Sitwala, MD, PhD - Medical Director
Form: MM RL1
PAGE 1 OF 1