

Example Client, XYZ123 1234 Warde Road Ann Arbor MI 48108 **EXAMPLE, REPORT** WX0000003039 M 12/05/1988 34 Y

		Molecula	r				
		Collected:	02/19/2022	06:06	Received:	02/19/202	2 06:06
Test Name	2	<u>Result</u>	Flag	Ref-Range	<u>s l</u>	<u>Jnits</u>	<u>Site</u>
Factor '	V Leiden Mutation Anal	lvsis					
Factor V Le	eiden Mutation	Heterozygous	AB				WMRL
	Both the wild type (WT) F5 gene and the pathogenic F5 Leiden variant (c.1691G>A) were detected indicating a heterozygous c.1691G>A genotype for this specimen. Heterozygosity for the c.1691G>A variant is associated with activated protein C resistance and an increased risk for venous thromboembolism (VTE). Factor V Leiden thrombophilia is inherited in an autosomal dominant manner and adults heterozygous for the F5 Leiden variant have a 3 to 8-fold increased risk of VTE. The prevalence of c.1691G>A heterozygosity is highest in individuals of European ancestry (5.2%). The variant is less common in Americans with Hispanic (2.2%), African						
	<pre>(1.2%), Asian (<0.5%), ancestry. The overall r individuals is approxim for both F5 Leiden and variant may experience tends to be more severe counseling is recommend testing asymptomatic famous of the second testing asymptomatic famous of testing asympt</pre>	and Native American (1. risk of a first VTE in he hately 0.5%. Patients he the G20210A (c.*97G>A) p earlier onset of thrombo than individual alleles led to help determine the mily members.	25%) eterozygou prothromk osis that s. Geneti e benefit	ous is vin c c c of			
	This test was performed (Roche) - an in vitro d real-time quantitative for the detection and g (F5) gene. The test det (WT) F5 gene and the pa known as the F5 Leiden from whole blood specim with suspected thrombop the cobas z 480 analyze amplification and detect this test is 0.1 ng/uL reaction).	a using the cobas® Factor liagnostic device that us Polymerase Chain Reaction genotyping of the human sects the presence of the thogenic c.1691G>A variation variant) in genomic DNA mens as an aid in diagnos whilia. The cobas® Factor er are used together for stion. The limit of deter of genomic DNA (2.5 ng/)	r V Test ses on (qPCR) factor V e wild ty ant (also isolated sing pati r V Test automate ction for PCR	rpe l ents and ed			
			Repo	rted Date:	02/19/2022	06:06 F	5LM
	Performing Site: WMRL: WARDE MEDICAL LABORATORY 300 West Textile Road Ann Arbor ML48108						

LAB: L - LOW, H - HIGH, AB - ABNORMAL, C - CRITICAL, . - NOT TESTED

Ordered By: CLIENT CLIENT WX0000000001595

Kajal V. Sitwala, MD, PhD - Medical Director Form: MM RL1 PAGE 1 OF 1