

Example Client, XYZ123 1234 Warde Road Ann Arbor MI 48108 EXAMPLE, REPORT W WX0000003827 M 07/08/1978 45 Y

		Referral Te	•				
		Collected	1: 08/30/2023	5 14:47	Received:	08/30/2023	14:47
<u>Test Name</u>		<u>Result</u>	<u>Flag</u>	Ref-Range	<u>s l</u>	<u>Jnits</u>	<u>Site</u>
Chlamydia Species AB Panel IgM							
Chlamydia IgM	Panel Interp	See Note					ARRL
INTERPRETIVE INFORMATION: Chlamydia IgM Panel							
gei may tha tis tra coi sar tha sar for and	The Chlamydia antibody test contains both species- and genus-specific antigens, and serological cross-reactions may be seen in both acute and convalescent samples (less than 1:128). A C. pneumoniae-specific reaction will exhibit titers twofold or greater than titers observed with C. trachomatis or C. psittaci serology. Ideally, acute and convalescent samples should be tested simultaneously at the same facility. If the sample submitted was collected during the acute-phase of illness, submit a marked convalescent sample within 25 days for paired testing. Seroconversion, a fourfold or greater rise in antibody titer between acute and convalescent sera, is considered strong evidence of recent infection.						
ps: tra	The Chlamydia microimmunofluorescent assay utilizes C. psittaci, C. pneumoniae, and nine serotypes of C. trachomatis. It does not include the LGV strains of C. trachomatis.						
de apj wa: in Pe: 500 Sa:	is test was developed an termined by ARUP Laborat proved by the US Food an s performed in a CLIA ce tended for clinical purp rformed By: ARUP Laborat 0 Chipeta Way lt Lake City, UT 84108 boratory Director: Jonat	ed or					
C. pneumoniae	e IgM Titer	<1:20		<1:20			ARRL
C. psittaci IgM ⁻	•	1:40	н	<1:20			ARRL
C. trachomatis IgM Titer		<1:20		<1:20			ARRL
Performing Site ARRL: ARUP REFERENCE LAB 500 Chipeta Way Salt Lake City UT 84108122							
			Reported			, 0	

LAB: L - LOW, H - HIGH, AB - ABNORMAL, C - CRITICAL, . - NOT TESTED

Ordered By: KAJAL SITWALA, MD, PhD WX0000000002365