



LABORATORY REPORT

Example Client, XYZ123
1234 Warde Road
Ann Arbor MI 48108

EXAMPLE, REPORT W
WX0000003827 M 07/08/1978 45 Y

Chemistry

Collected: 11/24/2023 08:55 Received: 11/24/2023 08:55

<u>Test Name</u>	<u>Result</u>	<u>Flag</u>	<u>Ref-Ranges</u>	<u>Units</u>	<u>Site</u>
Prostatic Specific Antigen (PSA), Diagnostic					
Prostate Specific Ag, Diagnostic	2.5		<=4.0	ng/mL	WMRL

After radical prostatectomy, the reference interval is less than 0.1 ng/mL if there is no residual disease. In healthy males without prostatectomy, the reference interval is 4.0 or less. The lower limit of detection is 0.1 ng/mL.

The intended use for this assay is measurement of serum PSA in conjunction with digital rectal examination as an aid in the detection of prostate cancer in men aged 50 years or older. There is also a second indication for this assay, which is serial measurement of PSA to aid in the prognosis and management of patients with prostate cancer; as such, specimens from patients under the age of 50 are not rejected by the laboratory. Interpretation of results, and use of provided reference range (established for the indication of prostate cancer detection in men 50 and older), should take intended uses into consideration.

This test was performed using the Beckman Coulter method, calibrated to the original Hybritech Tandem-R assay. PSA values obtained with other assay methods or kits cannot be used interchangeably with results obtained by the Beckman Tandem-R method.

Results cannot be interpreted as absolute evidence of the presence or absence of malignant disease.

Reported Date: 11/24/2023 08:56 PSADX

Performing Site:

WMRL: WARDE MEDICAL LABORATORY 300 West Textile Road Ann Arbor MI 48108

LAB: L - LOW, H - HIGH, AB - ABNORMAL, C - CRITICAL, . - NOT TESTED

F524000001
WX0000003827

Printed D&T: 11/24/23 08:56

Ordered By: KAJAL SITWALA, MD, PhD
WX00000000002365

Kajal V. Sitwala, MD, PhD - Medical Director

Form: MM RL1

PAGE 1 OF 1