



LABORATORY REPORT

QC ACCOUNT (WARDE)
300 W. TEXTILE
ANN ARBOR MI 48108

EXAMPLE, REPORT
WX0000000158 M 07/08/1968 57 Y

Referral Testing

Collected: 01/16/2026 12:09 Received: 01/16/2026 12:09

Table with 6 columns: Test Name, Result, Flag, Ref-Ranges, Units, Site. Rows include Babesia microti Antibodies (IgG, IgM) w/ Reflex to Titers.

Confirmation with a blood smear or PCR is recommended for diagnosis of acute Babesiosis. A single acute antibody titer is not sufficient to establish a diagnosis. IgG titers > or = 1:1024 or presence of IgM suggest recent infection. Antibodies may persist for months to years after clearance of infection. The extent of cross-reactivity between Babesia species is variable and other species (e.g., Babesia duncani) may not be detected by this assay. This test was developed and its analytical performance characteristics have been determined by Quest Diagnostics. It has not been cleared or approved by the FDA. This assay has been validated pursuant to the CLIA regulations and is used for clinical purposes. Test Performed by Quest, Chantilly, Quest Diagnostics Nichols Institute, 14225 Newbrook Drive, Chantilly, VA 20151 Patrick W Mason, M.D., Ph.D., Director of Laboratories (703) 802-6900, CLIA 49D0221801

Table with 6 columns: Test Name, Result, Flag, Ref-Ranges, Units, Site. Rows include Babesia microti Ab, IgG, Titer and Babesia microti Ab, IgM, Titer.

Reported Date: 01/16/2026 12:10 BMABT

LAB: L - LOW, H - HIGH, AB - ABNORMAL, C - CRITICAL, . - NOT TESTED